

Europäisches Patentamt

European Patent Office

Office européen des brevets



1) Publication number:

0 503 440 A1

(2)

## EUROPEAN PATENT APPLICATION

- 21 Application number: 92103592.9
- (1) Int. Cl.5: A61K 31/40, A61K 9/28

- 2 Date of filing: 02.03.92
- 3 Priority: 08.03.91 GB 9104890
- Date of publication of application:
   16.09.92 Bulletin 92/38
- Designated Contracting States:
   AT DE DK ES GR LU MC NL PT SE
- 7) Applicant: GLAXO GROUP LIMITED Glaxo House, Berkeley Avenue Greenford, Middlesex UB6 0NN(GB)
- ② Inventor: Padfield, John Malcolm
  Glaxo Manufact. Serv. Ltd, Stockley Park
  West
  Uxbridge, Middlesex UB11 1BU(GB)
  Inventor: Winterborn, Ian Keith, Glaxo Group
  Research Ltd.
  Park Road
  Ware, Hertfordshire SG12 0DG(GB)
  Inventor: Phillips, Anthony John, Glaxo Group
  Research Ltd.
  Park Road
- Representative: Filler, Wendy Anne et al Glaxo Holdings p.l.c., Glaxo House, Berkeley Avenue Greenford, Middlesex UB6 0NN(GB)

Ware, Hertfordshire SG12 0DG(GB)

(9) Compositions containing sumatriptan.

A pharmaceutical composition for oral adminstration comprising a film-coated solid dosage form including 3[2-(dimethylamino)ethyl]-N-methyl-1H-indole-5-methanesulphonamide or a pharmaceutically acceptable salt or solvate thereof as active ingredient.

The film-coated solid dosage forms are of use in the treatment of conditions associated with cephalic pain, in particular migraine.

EP 0 503 440 A1

The present invention relates to a pharmaceutical composition containing as active ingredient 3-[2-(dimethylamino)ethyl]-N-methyl-1H-indole-5-methanesulphonamide, in particular a composition for oral administration.

3-[2-(dimethylamino)ethyl]-N-methyl-1H-indole-5-methanesulphonamide, which may be represented by the formula (I)

20

and its physiologically acceptable salts and solvates are disclosed in UK Patent Specification No. 2162522. The compound of formula (I) exhibits selective vasoconstrictor activity and is useful in the treatment of migraine.

Oral administration constitutes the generally preferred route for administration of pharmaceuticals since this route is particularly convenient and acceptable to patients. Unfortunately oral compositions may be associated with certain disadvantages in the treatment of conditions associated with cephalic pain. For example, such conditions, particularly migraine are associated with gastrointestinal dysfunction in the form of delayed gastric emptying. This leads to both a delay and an impairment of drug absorption and it is generally accepted that oral formulations of drugs for the treatment of such conditions should be administered in the form of a liquid preparation.

Numerous clinical studies have demonstrated the effectiveness of the compound of formula (I) in migraineurs. Hitherto, the drug has always been administered either by parenteral injection or in the form of a dispersible tablet which is dispersed in drinking water prior to oral administration. This mode of oral administration was believed to minimise the potential problems associated with gastrointestinal dysfunction in migraineurs.

However, it has been found that the compound of formula (I) has a particularly unpleasant taste. When the compound of formula (I) is administered orally this unpleasant taste may exacerbate the nausea and vomiting associated with migraine.

The present invention provides a particularly advantageous formulation suitable for oral administration of the compound of formula (I).

There is thus provided according to the invention a pharmaceutical composition for oral administration comprising a film-coated solid dosage form including 3-[2-(dimethylamino)ethyl]-N-methyl-1H-indole-5-methanesulphonamide or a pharmaceutically acceptable salt or solvate thereof as active ingredient.

As used herein the term "film-coated solid dosage form" means a solid core comprising the active ingredient, which solid core is substantially covered with a film coating.

The compositions of the invention may comprise, for example, granules, tablets or capsules. Preferably the compositions of the invention will comprise tablets, most preferably compressed tablets.

There is provided in a preferred aspect of the invention a film coated tablet comprising a tablet core containing an effective amount of 3-[2-(dimethylamino)ethyl]-N-methyl-1H-indole-5-methanesulphonamide or a pharmaceutically acceptable salt or solvate thereof as active ingredient and a film coat on the tablet core.

We have found that the unpleasant taste associated with oral administration of the compound of formula (I) is substantially eliminated by the formulations of the present invention. The film coating also makes the formulations easier to handle and reduces potentially hazardous dust formation occurring during the packaging or administration of the drug. Surprisingly these advantages are attained without any significant loss in the bioavailability of the compound of formula (I) when compared to aqueous solutions or dispersible tablet formulations for oral administration to migraineurs. Film-coated tablets according to the invention are therefore surprisingly effective in the treatment of migraine.

It is preferred that 3-[2-(dimethylamino)ethyl]-N-methyl-1H-indole-5-methanesulphonamide should be employed in the compositions of the invention in the form of a physiologically acceptable salt. Such salts include salts of inorganic or organic acids such as hydrochloride, hydrobromide, sulphate, nitrate, phosphate, formate, mesylate, citrate, benzoate, fumarate, maleate, tartrate and succinate salts. Most preferably

3-[2-(dimethylamino)ethyl]-N-methyl-1H-indole-5-methanesulphonamide will be employed in the compositions of the invention in the form of its succinate (1:1) salt.

The film coating comprises a polymer. Suitable polymers include cellulose ethers, for example, hydroxypropyl methylcellulose, hydroxypropyl cellulose or methylcellulose, and copolymers of methacrylic acid and methyl methacrylate.

Preferably the film coating will comprise hydroxypropylmethyl cellulose.

The total film coating solids are generally applied to the solid dosage form, for example the tablet core, in an amount of from 2 to 5% w/w, preferably from 3 to 4% w/w, based on the weight of the solid dosage form.

The film coating may additionally comprise any pharmaceutically acceptable colourants or opacifiers including water soluble dyes, aluminium lakes of water soluble dyes and inorganic pigments such as titanium dioxide and iron oxide. Suitable colourants or opacifiers may comprise from 5% to 65% w/w, preferably from 25 to 50% w/w, based on the dry weight of film coating.

The film coating may also contain one or more plasticizing agents conventionally used in polymeric film coatings, for example polyethylene glycol, propylene glycol, dibutyl sebecate, mineral oil, sesame oil, diethyl phthalate and triacetin. Suitable plasticizing agents may comprise 1 to 40% preferably 5 to 20% w/w based on the dry weight of the film coating.

In addition to the compound of formula (I) or a physiologically acceptable salt or solvate thereof, compositions of the invention will preferably comprise pharmaceutically acceptable carriers and excipients, such as binding agents (e.g. pregelatinised maize starch, polyvinylpyrrolidone or hydroxypropylmethylcellulose); fillers (e.g. lactose, sucrose, mannitol, maize starch, microcrystalline cellulose or calcium hydrogen phosphate); lubricants (e.g. stearic acid, polyethylene glycol, magnesium stearate, talc or silica); disintegrants (e.g. potato starch, sodium starch glycollate or croscarmellose sodium); or wetting agents (e.g. sodium lauryl sulphate).

For the preparation of compositions according to the invention 3-[2-(dimethylamino)ethyl]-N-methyl-1H-indole-5-methanesulphonamideor a physiologically acceptable salt or solvate thereof may be blended with suitable excipients and, if desired, granulated. Preferably 3-[2-(dimethylamino)ethyl]-N-methyl-1H-indole-5-methanesulphonamide will be granulated with a filler before admixture with the other excipients. Most preferably the filler employed will be lactose. Tablets in uncoated form may be prepared, for example, by compression of the powder blend or granulate, using a lubricant as an aid to tabletting. Compressed tablets are preferred.

The solid dosage form is then film-coated using a suspension comprising a suitable polymer in a suitable solvent. The preferred solvent for the film coating components is purified water but various classes of organic solvents commonly used in this art such as alcohols, ketones, ethers and chlorinated hydrocarbons, for example ethanol, acetone, methylene chloride and the like, may also be used. The solvent does not appear in the final product. The amount of solvent may be varied according to the equipment and coating conditions used to produce an aesthetically coated tablet.

The amount of 3-[2-(dimethylamino)ethyl]-N-methyl-1H-indole-5-methanesulphonamide, preferably in the form of a physiologically acceptable salt, employed in the compositions of the invention will preferably be in the range of about 25mg to about 200mg, most preferably about 50mg or 100mg, expressed as the weight of free base.

A further aspect of the invention provides a method of treating a mammal, including man, suffering from or susceptible to conditions associated with cephalic pain such as cluster headache, chronic paroxysmal hemicrania, headache associated with vascular disorders, headache associated with substances or their withdrawal (for example drug withdrawal), tension headache and in particular migraine which comprises oral administration of a pharmaceutical composition comprising a film-coated solid dosage form of 3-[2-dimethylamino)ethyl]-N-methyl-1H-indole-5-methanesulphonamide or a pharmaceutically acceptable salt or solvate thereof as active ingredient. It will be appreciated that reference to treatment is intended to include prophylaxis as well as the alleviation of established symptoms.

It will be appreciated that the precise therapeutic dose of the active ingredient will depend on the age and condition of the patient and the nature of the condition to be treated and will be at the ultimate discretion of the attendant physician.

However, in general effective doses for the treatment of conditions associated with cephalic pain, for example acute treatment of migraine, will lie in the range of 10 to 500mg, preferably 20 to 300mg, most preferably 25 to 200mg, for example 50mg or 100mg of the active ingredient per unit dose which could be administered in single or divided doses, for example, 1 to 4 times per day.

The invention is further illustrated by the following non-limiting examples wherein the active ingredient is 3-[2-(dimethylamino)ethyl]-N-methyl-1H-indole-5-methanesulphonamide (1:1) succinate.

## Example 1

25

30

35

40

50

# Tablet cores

	140.01.00	
5		Unit formula (mg/tablet)
10	Active ingredient/lactose granule *	280.0
	Microcrystalline Cellulose Ph Eur	15.5
15	Croscarmellose Sodium USNF	3.0
	Magnesium Stearate Ph Eur	1.25 - 1.75
20	*Active ingredient/lactose granule	

Compound of formula (I) succinate 140.0\*\*

Lactose Ph Eur 170 mesh 140.0

Purified water Ph Eur qs +

+ The water does not appear in the final product. Typical range 100-140g per kg of blend

\*\* Equivalent to 100mg free base

Coating Suspension		
	% w/w	
Hydroxypropyl methylcellulose Ph Eur	10.0	
Opaspray white #	5.0	
Purified Water Ph Eur to	100.0++	

+ + The water does not appear in the final product. The maximum theoretical weight of solids applied during coating is 11mg/tablet. # Opaspray white is a proprietory film coating suspension, obtainable from Colorcon Ltd, UK, which contains hydroxypropyl methylcellulose and titanium dioxide.

The active ingredient and lactose were mixed together and granulated by the addition of purified water. The granules obtained after mixing were dried and passed through a screen, and the resulting granules were then mixed with the other tablet core excipients. The mix was compressed into tablets. The tablets were then film coated using the coating suspension in conventional film coating equipment.

## Example 2

The tablet cores were prepared as described in Example 1. The tablets were then film coated using the coating suspension given below and conventional film coating equipment.

Coating Suspension	
	% w/w
Opadry pink## Purified water Ph. Eur. to	5.3 100.0 + +

+ + The water does not appear in the final product. The maximum theoretical weight of solids applied during coating is 9mg/tablet.

## Opadry pink is a proprietory film coating material, obtainable from Colorcon Ltd, UK which contains hydroxypropyl methylcellulose, titanium dioxide, red iron oxide and triacetin.

#### Claims

5

10

15

- A pharmaceutical composition for oral administration comprising a film-coated solid dosage form including 3-[2-(dimethylamino)ethyl]-N-methyl-1H-indole-5-methanesulphonamide or a pharmaceutically acceptable salt or solvate thereof as active ingredient.
- 2. A pharmaceutical composition as claimed in Claim 1 wherein the active ingredient is in the form of its succinate (1:1) salt.
  - 3. A pharmaceutical composition as claimed in Claim 1 or Claim 2 in the form of a tablet.
- 4. A pharmaceutical composition as claimed in Claim 3 in the form of a compression tablet.
  - 5. A pharmaceutical composition as claimed in any one of Claims 1 to 4 wherein the film coating comprises a polymer.
- 6. A pharmaceutical composition as claimed in Claim 5 wherein the polymer is hydroxypropyl methylcellulose.
  - A pharmaceutical composition as claimed in any one of Claims 1 to 6 wherein the film coating comprises 2 to 5% w/w based on the weight of the solid dosage form.
  - A pharmaceutical composition as claimed in any one of Claims 1 to 7 which comprises 25 to 200mg of active ingredient.
- 9. A method of treating a mammal, including man, suffering from or susceptible to conditions associated with cephalic pain such as cluster headache, chronic paroxysmal hemicrania, headache associated with vascular disorders, headache associated with substances or their withdrawal (for example drug withdrawal), tension headache and in particular migraine which comprises oral administration of a pharmaceutical composition comprising a film-coated solid dosage form of 3-[2-dimethylamino)ethyl]-N-methyl-1H-indole-5-methanesulphonamide or a pharmaceutically acceptable salt or solvate thereof as active ingredient.
  - 10. A process for the preparation of a pharmaceutical composition as claimed in any one of Claims 1 to 8 which comprises applying a film coating to a solid dosage form of the active ingredient by a film coating technique.

55



# PARTIAL EUROPEAN SEARCH REPORT

Application Number

which under Rule 45 of the European Patent Convention shall be considered, for the purposes of subsequent proceedings, as the European search report

EP 92 10 3592

				1
]	DOCUMENTS CONSI			
Category	Citation of document with it of relevant parts	edication, where exprepriate, sanges	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int. Cl. 5)
D,X	GB-A-2 162 522 (GL * Claims; page 10,	AXO) lines 2-4 *	1-10	A 61 K 31/40 A 61 K 9/28
A	EP-A-0 147 107 (GL * Claims 1,8; page	AXO) 37, lines 8-10 *	1-10	
				TECHNICAL FIELDS SEARCHED (Int. CL5)  A 61 K
INCO	MPLETE SEARCH			1
Chairns se Chairns se Chairns se Chairns se Chairns se Reason fo Remain method their cari	ch Division considers that the present items of the European Patent Convent inlagful search into the state of the auxorbed completely: carched incompletely: or the limitation of the search: ark: Although cland of treatment rapy (Art. 52(4) ried out and base the composition.	taim 9 is directed of the human book EPC) the search	ed to a ly by has been	
	Photo of search	Date of completion of the s		RPONI U.
THI	E HAGUE	14-04-1992		
X : par Y : par doc A : tec O : no	CATEGORY OF CITED DOCUME ricularly relevant if taken alone ricularly relevant if combined with an ument of the same category shoological background a-written disclosure comediate document	E : earlier ; after th  other D : docume L : docume	r principle underlying the satest document, but pulle filing date nt cited in the application of the same patent fam nt	on s